

# Photobiomodulation: Deep Tissue Laser Therapy

Brian A. Pryor, PhD, Luis H. De Taboada, MSEE, Wendy S. Frydrych, PhD

## Introduction

Deep Tissue Laser Therapy uses a process called photobiomodulation to change the condition of damaged tissue by stimulating cellular metabolism, thereby accelerating the healing process. As light pours into the tissue, photons will be scattered, reflected and absorbed. Lasers operating in the near-infrared spectrum from 650 to 1300 nanometers can penetrate to deep tissue structures. Light that penetrates into the tissue can be absorbed by melanin, hemoglobin, oxyhemoglobin, and water. Energy from these absorption events will be dissipated as heat, generating a soothing warmth in the tissue.

The primary target for photobiomodulation is the cytochrome c complex which is found in the inner membrane of the mitochondria. cytochrome c is a vital component of the electron transport chain that drives cellular metabolism. As light is absorbed, cytochrome c is stimulated, leading to increased production of adenosine triphosphate (ATP), the molecule that facilitates energy transfer within the cell. In addition to ATP, laser stimulation also produces free nitric oxide and reactive oxygen species. Nitric oxide is a powerful vasodilator and an

important cellular signaling molecule involved in many physiological processes. Reactive oxygen species have been shown to affect many important physiological signaling pathways including the inflammatory response. In concert, the production of these signaling molecules has been shown to induce growth factor production, to increase cell proliferation and motility, and to promote extracellular matrix deposition and pro-survival pathways. Outside the cell, nitric oxide signaling drives vasodilation which improves microcirculation in the damaged tissue, delivering oxygen, vital sugars, proteins, and salts while removing wastes.

The recent development of higher-power, Class IV systems affords the clinician the ability to efficiently deliver adequate doses of light deep into tissue to reduce pain, reduce inflammation, and accelerate healing. Additionally, a contact treatment applicator that can be used to compress superficial tissues, displacing excess fluid, also enhances laser penetration to deep structures.

---

## Clinical Applications & Endorsements

Laser Therapy is endorsed by professional clinical organizations, including the World Health Organization (WHO), American Physical Therapy Association (APTA) and the International Association for the Study of Pain (IASP®). Over 3,000 research studies have been conducted in the field. Deep Tissue Laser Therapy can be used to treat pain and inflammation associated with the following conditions:

- Bursitis
- Disc Issues
- Epicondylitis
- Low Back Pain
- Neck Pain
- Neuropathy
- Plantar Fasciitis
- Sciatica
- Shoulder & Knee
- Sports Injuries
- Sprains & Strains
- Tendonitis & Tendinosis

---

## References

- Anders JJ, Moges H, Wu X, Erbele ID, Alberico SL, Saidu EK, Smith JT, Pryor BA. In vitro and in vivo optimization of infrared laser treatment for injured peripheral nerves. *Lasers in Surgery and Medicine*. 2014; 46(1): 34-45. doi: 10.1002/lsm.22212
- Garcia CR, Martin RL, Houck J, Wukich DK. Achilles Pain, Stiffness, and Muscle Power Deficits: Achilles Tendinitis--Clinical Practice Guidelines Linked to the International Classification of Functioning, Disability, and Health from the Orthopaedic Section of the American Physical Therapy Association. *J Orthop Sports Phys Ther*. 2010; 40.9: A1-A26. doi:10.2519/jospt.2010.0305
- Cotler HB, Chow RT, Hamblin MR, Carroll J. The Use of Low Level Laser Therapy (LLLT) For Musculoskeletal Pain. *MOJ Orthop Rheumatol*. 2015; 2(5): 00068. doi: 10.15406/mojor.2015.02.00068
- Haldeman S, Carroll L, Cassidy D, Schubert J, Nygren Å. The bone and joint decade 2000–2010 task force on neck pain and its associated disorders. *Spine*. 2008; 33(4S): S5-S7.
- International Association for the Study of Pain. "Myofascial Pain." Global Year Against Musculoskeletal Pain. IASP. October 2009-October 2010. Web. August 4, 2012. < <http://www.iasp-pain.org/Content/NavigationMenu/GlobalYearAgainstPain/20092010MusculoskeletalPain/FactSheets/default.htm>>.
- Karlekar A, Bharati S, Saxena R, Mehta K. Assessment of feasibility and efficacy of class IV laser therapy for postoperative pain relief in off-pump coronary artery bypass surgery patients: A pilot study. *Annals of Cardiac Anaesthesia*. 2015; 18(3): 317-322. doi: 10.4103/0971-9784.159800
- Knapp DJ. Postherpetic neuralgia: case study of class 4 laser therapy intervention. *The Clinical Journal of Pain*. 2013; 49(10): e6-e9. doi: 10.1097/AJP.0b013e31828b8ef8
- Nguyen LM-D, Malamo AG, Larkin-Kaiser KA, Borsa PA, Adhietty PJ. Effect of near-infrared light exposure on mitochondrial signaling in C2C12 muscle cells. *Mitochondrion*. 2014; 14: 42-48. doi: 10.1016/j.mito.2013.11.001
- Panton L, Simonavice E, Williams K, Majock C, Kim J-S, Kingsley JD, McMillan V, Mathis R. Effects of class IV laser therapy on fibromyalgia impact and function in women with fibromyalgia. *The Journal of Alternative and Complementary Medicine*. 2013. 19(5): 445-452. doi: 10.1089/acm.2011.0398
- Roberts DB, Kruse RJ, Stoll SF. The effectiveness of therapeutic class IV (10 W) laser treatment for epicondylitis. *Lasers in Surgery and Medicine*. 2013; 45(5): 311-317. doi: 10.1002/lsm.22140
- Tumilty S, Mani R, Baxter GD. Photobiomodulation and eccentric exercise for Achilles tendinopathy: a randomized controlled trial. *Lasers in Medical Science*. 2016; 31(1):127-135. doi: 10.1007/s10103-015-1840-4